

# LBAM Environmental Advisory Task Force Meeting 2/21/08

## **Biological Control of the Light Brown Apple Moth**

[Augmentation using egg parasitoids]

William Roltsch, Ph.D., Calif. Dept. of Food & Agriculture,  
Integrated Pest Control Branch, Biological Control Program

# Biological Control by Augmentation

- **Intent** - Utilize large numbers of *Trichogramma* to knock down localized, high-density populations of LBAM.
- **Background**
  - *T. pretiosum* and *T. platneri* are native to Calif.
  - *T. platneri* – 60% reduction of codling moth in walnut & pome fruit (Mills et al. 2000)



# LBAM Biology: life stages

Approx. 60 Eggs

Young &  
Late Larvae

Pre-pupa  
& pupa



Male



Female






















# Dark eggs parasitized by Trichogramma



LDAM egg parasitism study  
Trichogramma pretiosum  
13 Feb '08





# Egg Parasitism Study Outline

## Start:

Weekly parasite shipment received



## Parasite prep:

Held to complete development and emerge the day of study. Varied Temperatures.



## Test setup:

Parasites isolate in gel caps  
Sex is determined for each  
Honey applied to culture tube  
LBAM eggs selected  
Parasites paired into a single gel cap  
Parasites placed in culture tube w/eggs and plugged

## Holding Environment:

25 degrees C  
60-75% RH  
14:13 L:D



## Data collection:

24 & 48 hr. parasite survivorship  
10 day egg fate: egg hatch, parasite emergence



# Test of LBAM Eggs Attached by Two Trichogramma Species

Species & test date	Number of egg masses attacked	Mean percent eggs attacked per egg mass	LBAM egg hatch [unparasitized egg masses]
T. Platneri 7 Feb. 2008	5 of 34 egg masses	38% [range: 3-58%]	65%
T. pretiosum 13 Feb. 2008	9 of 20 egg masses	61% [range: 29-96%]	36% [Est. minimum]
T. Platneri 13 Feb. 2008	8 of 20 egg masses		

Have a great day